

**WELLESLEY PUBLIC SCHOOLS
ALLERGY EMERGENCY CARE PLAN (AECP)**



NAME _____ GRADE/TEACHER _____ SCHOOL YEAR _____

ALLERGY TO _____

DESCRIBE PREVIOUS ALLERGIC REACTION _____

DESCRIBE PREVIOUS ANAPHYLACTIC REACTION _____

USES INHALER NO ___ YES ___ DESCRIBE _____

*******SIGNS OF AN ALLERGIC REACTION*******

SYSTEM	SYMPTOMS (the severity of symptoms can quickly change)
*Mouth	Itching and swelling of the lips, tongue or mouth
*Throat	Itching and/or sense of tightness in the throat, hoarseness and hacking cough
*Skin	Hives, itchy rash and/or swelling about the face or extremities
*Stomach	Nausea, abdominal cramps, vomiting and/or diarrhea
*Lungs	Shortness of breath, repetitive coughing and/or wheezing
*Heart	"Thready pulse" and/or passing out

ACTION

If student is showing any signs or symptoms of an allergic reaction or if ingestion is suspected:

1. Contact nurse immediately. The nurse will administer Benadryl and/or EpiPen_____, or EpiPenJr._____.
2. If no nurse available (ie on field trip) the trained staff member will administer EpiPen_____, or EpiPen Jr._____.
3. Call 911, notify EMS of allergic reaction and EpiPen administration
4. Notify Parent/Guardian or Emergency Contact (if parent/guardian not available):

Name _____ Name _____ Name _____

Relation _____ Relation _____ Relation _____

Phone(H) _____ Phone(H) _____ Phone(H) _____

Phone(C) _____ Phone(C) _____ Phone(C) _____

Phone(W) _____ Phone(W) _____ Phone(W) _____

DO NOT HESITATE TO ADMINISTER MEDICATION OR CALL 911


Parent
Signature _____

EpiPen® and EpiPen® Jr. Directions

- Pull off gray activation cap.



- Hold black tip near outer thigh (always apply to thigh).



- Swing and jab firmly into outer thigh until Auto-Injector mechanism functions. Hold in place and count to 10. Remove the EpiPen® unit and massage the injection area for 10 seconds.

Twinject™ 0.3 mg and Twinject™ 0.15 mg Directions



- Pull off green end cap, then red end cap.
- Put gray cap against outer thigh, press down firmly until needle penetrates. Hold for 10 seconds, then remove.



SECOND DOSE ADMINISTRATION:

If symptoms don't improve after 10 minutes, administer second dose:

- Unscrew gray cap and pull syringe from barrel by holding blue collar at needle base.
- Slide yellow or orange collar off plunger.
- Put needle into thigh through skin, push plunger down all the way, and remove.



DIRECTIONS FOR USE ARE ALSO ON EPIPEN